



Diurnal use of space by captive adult broad-snouted caiman (*Caiman latirostris*): Implications for pen design

Luciano M. Verdade^{a,*}, Carlos I. Piña^{a,b}, Janaina L.O. Araújo^a

^aAnimal Ecology Lab, University of São Paulo, PO Box 09, Piracicaba, SP 13418-900, Brazil

^bProyecto Yacaré, CICyTTP-CONICET, Dr. Matteri y España, CP 3105, Diamante, Entre Ríos, Argentina

Received 4 February 2004; received in revised form 13 June 2005; accepted 15 June 2005

Abstract

Crocodylians generally present low reproductive success in captivity. The reasons for this are still unclear, but the faulty design of reproductive facilities can be a cause of stress, injuries and social disruption. This study explored the use of space by captive adult broad-snouted caiman in order to improve pen design for farming species. Caimans showed a predominant use of the pool and its margin during daytime. Even in our small (9 × 10 m) pens, caimans used pools and their margins significantly more than areas farther removed from pools. This pattern suggests that caiman farmers should maximize pool area and perimeter in pen design.

© 2005 Elsevier B.V. All rights reserved.

Keywords: Crocodylian husbandry; Caiman farm; Pen-design; Behavior

Crocodylians exhibit elaborate social behavior including a complex social structure, parental care, and a sophisticated communication system of vocalizations, displays, and scent marking (Ayarzaguena, 1983; Lang, 1987; Vliet, 1989). Some crocodylian species maintain individual territories (sensu Kaufmann, 1983) year-round. In other species, territoriality is largely restricted to the mating season, and females may establish a social hierarchy within a male's terri-

tory (Lang, 1987). Such a seasonal pattern is typical of temperate species such as the American alligator (*Alligator mississippiensis*) and the broad-snouted caiman (*Caiman latirostris*) (Lance, 2003; Piña et al., 2004). The use of space may be related to the individual social status (Grant, 1973). In general, aggressive and big individuals tend to dominate mating activities and habitat selection for nesting, foraging and basking (Gould and Gould, 1989).

Agonistic interactions can represent up to 15% of the total mortality in captivity for *C. latirostris* since in 80% of the fights at least one individual dies (Verdade, 1992). Under farming conditions, the broad-snouted caiman seems eventually to establish

* Corresponding author.

E-mail addresses: lmv@esalq.usp.br (L.M. Verdade), cidcarlos@infoaire.com.ar (C.I. Piña).

a social structure that decreases the occurrence of agonistic interactions (Verdade, 1992, 1995, 1999).

Because it is difficult to study crocodylian social behavior in the wild, many behavioral studies have been carried out in captivity (Lang, 1987; Vliet, 1989; Verdade, 1999; Piffer and Verdade, 2002; Piña, 2002). Such studies can show caiman farmers how to reduce negative effects of artificial environments on the welfare of captive crocodylians (Deag, 1981; Warwick et al., 1995).

The broad-snouted caiman is beginning to be commercially produced in Argentina and Brazil (Verdade and Piña, in press). In some provinces of Argentina, where healthy wild populations can still be found, sustained-yield exploitation is based on egg collection in wild and captive rearing of young animals (Larriera, 1990, 1992, 1993, 1994, 1995, 1998). In Brazil, *C. latirostris* has a larger geographical distribution, but its wild populations are generally smaller and more fragmented (Verdade, 1997; Verdade et al., 2002). Therefore, the establishment of a ranching program in Brazil is still unfeasible and the commercial management of the species requires captive maintenance of reproductive adults (i.e., “farming”, according to Hutton and Webb, 1992).

Adult crocodylians generally exhibit a low reproductive rate in captivity (Cardeilhac, 1989, 1990). The reasons for this are still unclear, but the configuration of reproductive facilities can be a major cause of stress, injuries and social disruptions (Arena and Warwick, 1995; Huchzermeyer, 2003). This study aims to determine the use of space by captive adult broad-snouted caiman in order to improve pen design for farming the species.

1. Materials and methods

This study was carried out at the caiman facilities of the Animal Ecology Lab of the University of São Paulo, in Piracicaba, State of São Paulo, Brazil (22°42.557'S, 47°38.246'W). The captive colony of the species was established in 1987 (Verdade, 2001). The reproductive groups of this study were assembled in 1992 and transferred to the current facilities in 1998. The caimans groups were assembled based on parenthood among individuals recorded in a local Studbook (Verdade and Kassouf-Perina, 1993) but

no caiman was housed with an animal >1.2 times its body-mass (as suggested by Verdade, 1992). Since 1992, 30% to 60% of females have nested every year (average of approximately 40%).

We used four adult groups of one male and two to four females in four reproductive enclosures (Table 1). Each enclosure was 9 × 10 m; each included a 6 × 4 m pool and five 2 × 2 m nesting shelters. Six “microhabitats” (sensu Douglas, 1976) were defined as indicated in Fig. 1.

We used a scanning observation method (Lehner, 1996) every hour and registered each animal's location in the microhabitats above, during daytime (07:00–18:00). Observations were conducted from a 10 m high tower located 2 m outside the facility perimeter. Individuals were identified by their natural marks and tail notching marks. However relevant, nocturnal observations were not carried out in this study because of the impossibility of identifying the caimans in the dark. The use of lamps could affect caimans' behavior, as usually reported for crocodylians (Woodward and Marion, 1978; Hutton et al., 1987); therefore they have not been used.

The study period comprised 90 days evenly distributed over 30 weeks from 10 March to 13 November 2003, for a total of 990 h of observation. This

Table 1
Space use for each animal at each reproductive enclosure

Enclosure	Animal	Water	Margin	Reproduction	Land
1	♀1 (repro)	≈	++	--	≈
1	♀2 (non-repro)	≈	++	≈	≈
1	♂	≈	++	--	≈
2	♀1 (repro)	--	++	≈	≈
2	♀2 (non-repro)	--	++	--	≈
2	♀3 (non-repro)	--	++	≈	≈
2	♂	≈	++	--	≈
3	♀1 (repro)	≈	++	--	≈
3	♀2 (non-repro)	≈	++	--	≈
3	♂	≈	++	--	≈
4	♀1 (non-repro)	--	≈	--	++
4	♀2 (non-repro)	--	--	--	++
4	♀3 (non-repro)	--	++	--	≈
4	♀4 (non-repro)	--	≈	--	++
4	♂	≈	++	--	≈

++: Microhabitats used more than by chance. --: Microhabitats used less than by chance based on its area. ≈: Microhabitats used by chance. M is the male of each enclosure. Reproductive females (“repro”) laid eggs on the subsequent nesting season.




Fig. 1. Caiman reproductive facilities at Animal Ecology Lab, University of São Paulo. a) Nesting shelters (NS), as suggested by [Bustard, 1975](#); b) Nesting shelter margins (MgNS); c) Margin opposite to the gates (MgFar); d) Margin close to the gates (MgGate); e) Inner margin (MgWall); f) Pool margin (MgPool); and, g) Pool.

period covered the entire year, except the nesting season (late November to early March in this region; [Verdade, 1995](#)), when animals exhibit specific reproductive behaviors (e.g., mating and nest attendance) that are beyond the scope of the present study. The frequency of microhabitat use by males and females during the study period was compared by χ^2 test: expectation under H_0 was that use would be proportional to area.

2. Results

Reproduction occurred in three of the four enclosures. In all enclosures, there was a significant difference among individuals in relation to the use of microhabitats ($\chi^2 > 46.4$, $12 < df < 18$, $P < 0.001$); however, at ARN1, ARN2 and ARN3 the pools' margins (MgPool) were used more frequently than by chance alone ([Table 1](#)). In ANR4 (where there was no reproductive activity) the animals used the microhabitats in a different way, some used land more than by chance,

others used margins and all but the male used water less frequently than by chance ([Table 1](#)).

We found a significant difference in microhabitat use between males and females ($\chi^2 = 1040.177$, $df = 18$, $P < 0.001$). Males used the microhabitats differently among each other either considering the whole period of study ($\chi^2 = 59.94$, $df = 9$, $P < 0.001$) or the seasons separately ($23.14 < \chi^2 < 48.89$, $df = 9$, $P < 0.005$). However, considering the area of each microhabitat, the four males stayed relatively longer in the edge of the water, relatively shorter near the nesting shelters, and equally in the water and on the ground ([Table 1](#)). The females, on the other hand, presented different time ratios at different microhabitats ([Table 1](#)).

There is a significant difference between females from distinct facilities in relation to the use of microhabitats ($\chi^2 = 534.035$, $df = 5$, $P < 0.001$). Reproductive females used the microhabitats differently among each other either considering the whole period of study ($\chi^2 = 169.96$, $df = 6$, $P < 0.001$) or during three of the four seasons separately (i.e., summer, falls and winter; $10.13 < \chi^2 < 147.12$, $df = 9$, $P < 0.001$); during the




Fig. 2. Use of microhabitats (% of time) by adult captive broad-snouted caiman at the four reproductive facilities of the Animal Ecology Lab, University of São Paulo.

spring there was no difference among them ($\chi^2=10.13$, $df=6$, $P=0.1194$). Female 4 was the only one that stayed relatively shorter at the edge of the water (Table 1). In her group (ARN4) the animals presented the biggest variation in the use of microhabitats and did not breed (Fig. 2).

3. Discussion

This study was carried out during daylight time when crocodilians spend most of their time basking (Ayarzagüena, 1983). Although there is some general difference between males and females, an individual pattern of space use seems rather predominant. Individual behavioral patterns have been described in mammals (Box, 1990) and birds (Ligon and Ligon, 1991), and may also occur with crocodilians. This may be related to territoriality and should be investigated on future studies. Captive caimans tend to form stable reproductive groups (one male: some females) when animals' body sizes do not differ more than 20% from each other (Verdade, 1992).

Caimans showed a predominant use of the pool and its margin during daytime. Even on the small facilities of this study, the areas more distant from the water were used significantly less, especially those near the gate. This pattern is relevant to the design of caiman breeding facilities for which the pool perimeter should be enlarged as much as possible, but not necessarily increasing the total pen area. As the animals use the pool margin (with shallow water) more frequently than the ground around them, those margins should also be enlarged and the "dry" ground decreased. This can be easily done by enlarging the pool area and decreasing the ground around. Considering the current pen design, an increase in the pool margin from 75 to 100 cm would result in an increase of approximately 17% in the pool area (from 24 to 28.1 m²), and an increase of approximately 38% in the pool margin (from 10.9 to 15 m²). Cemented margins in platform (like a step) instead of ramp form are possibly more adequate as they allow the animals to bask in shallow water and also to exit the water without crawling across companions, which can result in bites or fights.

It is noteworthy that in this study reproductive females presented similar patterns of space use during the non-reproductive period. During the spring,

female American alligators store energy for the breeding season in the forthcoming summer (Lance, 2003). It is likely that caiman reproductive females exhibit stereotyped thermoregulatory behavior in order to optimize energy storage and egg production. As a matter of fact, captive adult reproductive broad-snouted caiman were warmer than non-reproductive females during the spring, 4 to 6 weeks prior to egg-laying (Bassetti, 2002).

Although the nesting shelters were not frequently used during this study, they are used extensively by females to nest in and thereby possibly avoiding agonistic interactions during that period (Verdade, 1995). Therefore, they should be kept in future designs of caiman reproductive facilities. However, their form and number should take into consideration the number of females per reproductive group as well as the logistics of egg collection and transportation during management.

As crocodilians generally exhibit nocturnal behavior quite distinct from diurnal behavior, future studies should focus on the use of space by caimans during nighttime. In such situation, observations might be significantly improved by the use of new technologies such as starlight cameras, as their costs become more accessible to researchers in developing countries.

Acknowledgments

This study was supported by FAPESP (proc. # 01/01495-3). Carlos I. Piña has a Post-Doctorate fellowship from FAPESP (proc. # 03/09120-7) and Janaina L.O. Araújo had an undergraduate training fellowship from FAPESP (proc. # 03/00190-2).

References

- Arena, P.C., Warwick, C., 1995. Miscellaneous factors affecting health and welfare. In: Warwick, C., Frye, F.L., Murphy, J.B. (Eds.), *Health and Welfare of Captive Reptiles*. Chapman & Hall, London, pp. 263–283.
- Ayarzagüena, J., 1983. Ecología del caimán de anteojos o baba (*Caiman crocodilus* L.) en los llanos de Apure (Venezuela). *Doñana Acta Vertebrata* 10 (3), 3–135.
- Bassetti, L.A., 2002. Comportamento de Termorregulação em Jacarés-dePapo-Amarelo (*Caiman latirostris*) Adultos em Cativeiro. Masters Dissertation. University of São Paulo. Piracicaba, SP, Brazil. 59 pp.

- Box, H.O., 1990. Responsiveness to environmental change: inter-relationships among parameters. In: Box, H.O. (Ed.), Primate Responses to Environmental Change. Chapman and Hall, London, pp. 57–89.
- Bustard, H.R., 1975. Captive breeding of crocodiles. In: Martin, R.D. (Ed.), Breeding Endangered Species in Captivity. Academic Press, London, pp. 43–47.
- Cardeilhac, P., 1989. Husbandry and preventative medicine practices that increase reproductive efficiency of breeding colonies of alligators. Aquaculture Market Develop Aid Program. 25 pp.
- Cardeilhac, P., 1990. Husbandry and preventative medicine practices that increase reproductive efficiency of breeding colonies of alligators (II). Aquaculture Market Development Aid Program. 47 pp.
- Deag, J.M., 1981. O comportamento social dos animais. EDUSP, São Paulo. 118 pp.
- Douglas, R.J., 1976. Spatial interactions and microhabitat selections of two locally sympatric voles. *Microtus montanus* and *Microtus pennsylvanicus*. *Ecology* 57, 346–352.
- Gould, J.L., Gould, C.G., 1989. Sexual Selection: Mate Choice and Courtship in Nature. Scientific American Library, New York. 277 pp.
- Grant, T.R., 1973. Dominance and association among members of a captive and a freeranging group of grey kangaroos (*Macropus giganteus*). *Animal Behaviour* 21 (3), 449–456.
- Huchzermeyer, F.W., 2003. Crocodiles Biology, Husbandry and Diseases. CABI Publishing, Oxon, U.K. 337pp.
- Hutton, J.M., Webb, G.J.W., 1992. An introduction to the farming of crocodylians. In: Luxmore, R.A. (Ed.), Directory of Crocodylian Farming Operations. IUCN—The World Conservation Union, Gland, Switzerland, pp. 1–39.
- Hutton, J.M., Loveridge, J.P., Blake, D.K., 1987. Capture methods for the Nile crocodile in Zimbabwe. In: Webb, G.J.W., Manolis, S.C., Whitehead, P.J. (Eds.), Wildlife Management: Crocodiles and Alligators. Surrey Beatty & Sons Pty Lim, Chipping Norton, Australia, pp. 243–247.
- Kaufmann, J.H., 1983. On the definition and functions of dominance and territoriality. *Biological Review* 58, 1–20.
- Lance, V.A., 2003. Alligator physiology and life history: the importance of temperature. *Experimental Gerontology* 38 (7), 801–805.
- Lang, J., 1987. Crocodylian behaviour: implications for management. In: Webb, G.J.W., Manolis, S.C., Whitehead, P.J. (Eds.), Wildlife Management: Crocodiles and Alligators. Surrey Beatty & Sons, Chipping Norton, Australia, pp. 273–294.
- Larriera, A., 1990. A program of monitoring and recovering of caimans populations in Argentina with the aim of management. Proceedings of the 10th working meeting of the Crocodile Specialist Group, of the Species Survival Commission of IUCN, Crocodiles, vol. 2. The World Conservation Union, Gland, Switzerland, pp. 1–5.
- Larriera, A., 1992. The experimental breeding station of *Caiman latirostris* at Santa Fe city. Proceedings of the 11th Working Meeting of the Crocodile Specialist Group of the Species Survival Commission of the IUCN, Crocodiles, vol. 1. The World Conservation Union, Gland, Switzerland, pp. 250–256.
- Larriera, A., 1993. The experimental breeding station of *Caiman latirostris* at Santa Fe city, Argentina. Zoocria de los Crocodylia Memorias de la I Reunión Regional del CSG, Grupo de Especialistas en Cocodrilos de la UICN. IUCN—The World Conservation Union, Gland, Switzerland, pp. 160–164.
- Larriera, A., 1994. *Caiman latirostris* ranching program in Santa Fe, Argentina, with the aim of management. Proceedings of the 12th Working Meeting of the Crocodile Specialist Group of the Species Survival Commission of the IUCN, Crocodiles The World Conservation Union, Gland, Switzerland, pp. 188–199.
- Larriera, A., 1995. Areas de nidificación y momento óptimo de cosecha de huevos de *Caiman latirostris* en Santa Fe, Argentina. In: Larriera, A., Verdade, L.M. (Eds.), La Conservación y el Manejo de Caimanes y Cocodrilos de América Latina, vol. 1. Fundación Banco Bica, Santo Tomé, Santa Fe, Argentina, pp. 221–232.
- Larriera, A., 1998. The *Caiman latirostris* ranching program in Santa Fe, Argentina. The first commercial rearing. Proceedings of the 14th Working Meeting of the Crocodile Specialist Group, Crocodiles IUCN—The World Conservation Union, Gland, Switzerland, pp. 379–385.
- Lehner, P.N., 1996. Handbook of Ethological Methods, 2nd ed. Cambridge University Press, Cambridge, U.K. 672 pp.
- Ligon, J.D., Ligon, S.H., 1991. The cooperative breeding behavior of the green woodhoopoe. In: Mock, D.W. (Ed.), Behavior and Evolution of Birds. W.H. Freeman and Company, New York, pp. 107–119.
- Piffer, T.R.O., Verdade, L.M., 2002. *Caiman latirostris* (broad-snouted caiman) courtship behavior. *Herpetological Review* 33 (2), 132–133.
- Piña, C.I., 2002. *Caiman latirostris* (Broad-snouted Caiman). Thermoregulation. *Herpetological Review* 33, 133.
- Piña, C.I., Larriera, A., Siroski, P., 2004. Cocodrilos en la region litoral: especies, distribución geográfica y modo de vida. *Insuego, Miscelanea* 12, 317–322.
- Verdade, L.M., 1992. Agonistic social behavior of broad-nosed Caiman (*Caiman latirostris*) in captivity: implications to reproductive management. Proc. 11th Working Meeting Crocodile Specialist group, Crocodiles, vol. 2. IUCN—The World Conservation Union, Gland, Switzerland, pp. 200–217.
- Verdade, L.M., 1995. Biología reproductiva do jacaré-de-papo-amarelo (*Caiman latirostris*) em São Paulo, Brasil. In: Larriera, A., Verdade, L.M. (Eds.), La Conservación y el Manejo de Caimanes y Cocodrilos de América Latina, vol. 1. Fundación Banco Bica, Santo Tomé, Santa Fé, Argentina, pp. 57–79.
- Verdade, L.M., 1997. Manejo e conservação do jacaré-de-papo-amarelo (*Caiman latirostris*) no Estado de São Paulo. In: Valladares-Padua, C.B., Bodmer, R.E., Cullen Jr., L. (Eds.), Manejo e Conservação de Vida Silvestre no Brasil. CNPq, Brasília, pp. 222–231.
- Verdade, L.M., 1999. *Caiman latirostris* (broad-snouted caiman) behavior. *Herpetological Review* 30 (10), 38–39.
- Verdade, L.M., 2001. O programa experimental de criação em cativeiro do jacaré-de-papo-amarelo (*Caiman latirostris*) da ESALQ/USP: histórico e perspectivas. In: Mattos, W.R.S. (Ed.), A Produção Animal na Visão dos Brasileiros. Sociedade Brasileira de Zootecnia, Piracicaba, SP Brasil, pp. 559–564.

- Verdade, L.M., Kassouf-Perina, 1993. Studbook Regional do Jacaré-de-Papo-Amarelo (*Caiman latirostris*)—1992/1993. Sociedade de Zoológicos do Brasil, Sorocaba. 46 pp.
- Verdade, L.M., Piña, C.I., in press. *Caiman latirostris*. Catalog of the Society for the Study of American Amphibians and Reptiles. 55 pp.
- Verdade, L.M., Zucoloto, R.B., Coutinho, L.L., 2002. Microgeographic variation in *Caiman latirostris*. Journal of experimental Zoology 294 (4), 387–396.
- Vliet, K.A., 1989. Social displays of the American alligator (*Alligator mississippiensis*). American Zoologist 29, 1019–1031.
- Warwick, C., Frye, F.L., Murphy, J.B., 1995. Health and Welfare of Captive Reptiles. Chapman and Hall. 299 pp.
- Woodward, A.R., Marion, W.R., 1978. An evaluation of factors affecting night-light counts of alligators. Proc. Conf. SEast Ass. Game Fish Wildl. Agencies, vol. 32, pp. 291–302.